



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SEVEN SPECTROSCOPIC BINARIES.

	Mag.	Spec.	R.A. 1910	Dec. 1910	Range in Velocity, km
Lalande 2682	7.8	K1	1 ^h 24 ^m .1	+ 21° 16'	— 17 to + 47
Boss 1082	5.3	G4	4 32 .8	+ 52 54	0 to — 66
A. G. C. 3881	8.0	A2p	9 24 .6	+ 33 42	Composite
Boss 2951	6.4	A1	11 2 .8	+ 23 49	— 13 to + 32
Boss 3108	5.8	K5	11 46 .4	— 4 50	— 2 to + 26
Boss 3284	6.7	A3p	12 30 .6	+ 18 52	Composite
Boss 5173	5.5	Aop	20 6 .8	+ 26 38	Composite

NOTES.

A. G. C. 3881.—Two plates show double lines. A photograph taken on January 5th gives for the velocities of the two components — 41 and + 81^{km}. Boss 3284. The third spectrogram showed the presence of two spectra. The velocities given by this plate are — 70 and + 57^{km}. Boss 5173.—The hydrogen lines in this star show great variations in width and intensity, due probably to the presence of two spectra. A relative displacement of about 150^{km} between the components is indicated by the first plate.

All of the photographs were taken with low dispersion.

WALTER S. ADAMS.

THREE STARS WITH GREAT RADIAL VELOCITIES.

The following stars have been found to have exceptionally high radial velocities:—

	Mag.	Spec.	R.A. 1910	Dec. 1910	μ	v	No. Plates
W. B. 3 ^h 617	7.2	F5	3 ^h 35 ^m .8	— 3° 31'	0".78	+ 114 ^{km}	3
W. B. 17 ^h 514	8.6	F1	17 30 .4	+ 6 4	0 .58	— 148	4
O. Arg. 20,452	8.1	Gop	20 18 .3	— 21 38	1 .21	— 179	4

WALTER S. ADAMS.

THE SPECTRUM OF T TAURI.

A recent photograph of the spectrum of the well-known irregular variable *T Tauri* shows some interesting peculiarities. The spectrum is compound and consists of a considerable number of well-defined bright lines superposed on a dark line spectrum of about type F5. The fact that bright lines cover *H* β , *H* γ , and H and K makes it difficult to classify the absorption spectrum closely. Measures of the photograph give the